



Burdekin

Variety	Q135 x QN61-1232
SEEDLING CODE	QA87-1413
APPEARANCE	Stalk green, yellow when exposed to sun Leaves dark green Open stool with many thin sticks Free trashing
FEATURES/COMMENTS	Growth cracks in stalks Suitable for average soils Low input
HARVESTING TIME	Early, Mid, Late
SUITED SOILS	
SEASONAL SUGAR -	
EARLY	Average
MID	Good Good
LATE	Good
YIELD	High
CCS	Moderate
FAST AND RELIABLE GERMINATION	Average
HARSH CONDITIONS	Average
LODGING TOLERANCE	Poor
TRASH YIELD	Unknown
WATERLOGGING	Poor
FLOOD	Unknown
WIND	Good
DISEASE REACTION -	
RESISTANT	Brown Rust, Chlorotic Streak, Leaf Scald,
INTERMEDIATE DECICEANT	Orange Rust, Red Rot
INTERMEDIATE - RESISTANT	Smut Posture atra PCD
INTERMEDIATE INTERMEDIATE - SUSCEPTIBLE	Pachymetra, RSD
SUSCEPTIBLE	Fiji Leaf Gall Lesion Nematode, Root Knot Nematode
UNKNOWN	Striate Mosaic, Yellow Leaf Virus
CANE GRUB TOLERANCE	Average
CROP MANAGEMENT PRACTICES	Reduce nitrogen rates to maximize CCS
CROI WANAGEMENT I RACTIGES	Ensure adequate drying down to maximize CCS
	Responsive to MODDUS®.
	If possible avoid early (June) harvest.
	Irrigation management needs to be tailored to suit the variety and soil type. Avoid overwatering on heavy soils.
	Do not dry down for too long when using for plants. 4 to 6 weeks is
	ideal.
	Charles attaids yours of and active for the time and to

Disclaimer

Copyright © 2013 • All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior permission of SRA. Disclaimer In this disclaimer a reference to "we", "us" or "our" means Sugar Research Australia Limited and our directors, officers, agents and employees. Although we do our best to present information that is correct and accurate, we make no warranties, guarantees or representations about the suitability, reliability, currency or accuracy of the information we present in this report for any purposes. Subject to any terms implied by law and which cannot be excluded, we accept no responsibility for any loss, damage, cost or expense incurred by you as a result of the use of, or reliance on, any materials and information appearing in this report. You, the user, accept sole responsibility and risk associated with the use and results of the information appearing in this report and you agree that we will not be liable for any loss or damage whatsoever (including through negligence) arising out of, or in connection with the use of this report. We recommend that you contact our staff before acting on any information provided in this report. Warning Our tests, inspections and recommendations should not be relied on without further, independent inquiries. They may not be accurate, complete or applicable for your particular needs for many reasons, including (for example) SRA being unaware of other matters relevant to individual crops, the analysis of unrepresentative samples or the influence of environmental, managerial or other factors on production.

Check outside rows of seed source for weathering cracks.

Do not plant late as germination may be unreliable.

